



Guidelines by stage

COTTON

STAGE	APPLICATION	BENEFITS
	Preplant <i>Soil Solution</i> incorporated @ 24 Oz/acre. 32 Oz/acre for no-till or overgrowing crops. Apply in the Fall or up to planting.	Composting and soil health.
	Seed Treatment <i>Seed Coat</i> @ 2 Oz (dry weight)/acre.	Final Stand.
	In-Furrow <i>PercPlus</i> @ 8 Oz/acre at planting.	Stress.
	1st True Leaf up to 4 Leaf <i>PercPlus</i> @ 16 Oz/acre foliar broadcast in 2 applications, 2 - 3 weeks apart. May be applied with glyphosate, other herbicides or insecticides.	Weather and stress.
	Side-Dress <i>Zinc Plus</i> @ 32 Oz/acre <i>Sulfur Plus</i> @ 32 Oz/acre <i>Potassium Plus</i> @ 32 Oz/acre. All are compatible with liquid nitrogen.	Supplemental.
	6th to 10th Node 3 applications of <i>Boron Plus</i> @ 4 Oz/acre at Pin-Head, Match-Head and First Bloom . Apply on 14 - 21 day intervals.	Foliar supplement.
	Heavy Boll Set <i>CropKarb</i> : Make two applications @ 32 Oz/acre each in the 4th & 6th or 5th & 7th week of heavy bloom & boll fill.	Maturity.

These DeltAg Guidelines have been utilized to improve overall crop health and are not intended to be utilized as a replacement for basic and customary fertilizer applications.

P.O. Box 4812

Greenville, MS 38704

Phone: 877-594-1500

www.seedcoat.com