



Guidelines for SOD

Dormancy Application: Apply *Soil Solution* @ 32 Oz/acre broadcast. This application can be made in the late Fall, Winter or early Spring, before turf starts to green up. May also be applied on green turf but not as beneficial for the specific season of application.

Initial Planting Seed Treatment: Apply *Seed Coat* @ 2 Oz (dry weight) per planted acre of seed.

Early Spring Application: Apply *PercPlus* @ 24 Oz/acre broadcast.

Late Spring Application: Apply *CropKarb* @ 32 Oz/acre applied with *Iron Plus* @ 8 Oz/acre.

Mid-Summer Application: Apply *CropKarb* @ 32 Oz/acre applied with *Iron Plus* @ 8 Oz/acre.

Pre-Harvest of Sod: Apply *PercPlus* @ 16 Oz/acre, preferably 2 to 3 weeks before harvest.

Post-Harvest: Apply *Soil Solution* @ 32 Oz/acre to the soil to enhance grow out after harvest.

NOTE: *CropKarb* may be replaced with a tank-mix of *Potassium Plus* @ 32 Oz, *PercPlus* @ 8 to 16 Oz, and *Boron Plus* @ 4 Oz. *Plant Power* may be substituted for *PercPlus* and vice-versa in any application and at the same rate.

Mixing: Always dilute *Potassium Plus* in water first, and follow with other ingredients in listed order to avoid excessive foaming.

These **DeltAg Guidelines** have been utilized to improve overall crop health and are not intended to be utilized as a replacement for basic and customary fertilizer applications. **“Other Crop Specific Guidelines are available upon request.”**